




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/82, 15/11, A01H 5/00, C08B 30/14		(11) International Publication Number: WO 95/26407
A1		(43) International Publication Date: 5 October 1995 (05.10.95)
(21) International Application Number: PCT/GB95/00634		Richard [GB/GB]; 10 Furness Close, Bedford, Bedfordshire MK41 8RN (GB). SIDEBOTTOM, Christopher, Michael [GB/GB]; 35 Waterloo Road, Bedford, Bedfordshire MK40 3PQ (GB). WESTCOTT, Roger, John [GB/GB]; 46 Castle Street, Wellingborough, Northamptonshire NN8 1LW (GB).
(22) International Filing Date: 22 March 1995 (22.03.95)		
(30) Priority Data: 9406022.5 25 March 1994 (25.03.94) GB 94305806.5 4 August 1994 (04.08.94) EP (34) Countries for which the regional or international application was filed: GB et al. 95300210.2 13 January 1995 (13.01.95) EP (34) Countries for which the regional or international application was filed: GB et al.		(74) Agent: KEITH W NASH & CO; Pearl Assurance House, 90-92 Regent Street, Cambridge CB2 1DP (GB).
(71) Applicant (for all designated States except US): NATIONAL STARCH AND CHEMICAL INVESTMENT HOLDING CORPORATION [US/US]; Investment Holding Corporation, Suite 27, 501 Silverside Road, Wilmington, DE 19809 (US).		(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LT, LU, LV, MD, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG).
(72) Inventors; and (75) Inventors/Applicants (for US only): COOKE, David [GB/GB]; 2 Hunts Path, Oakley, Bedfordshire MK43 7SR (GB). GIDLEY, Michael, John [GB/GB]; 45 Holmes Avenue, Raunds, Northamptonshire NN9 6SZ (GB). JOBLING, Stephen, Alan [GB/GB]; 19 Burwell Road, Eaton Socon, Huntingdon, Cambridgeshire PE19 3QQ (GB). SAFFORD,		Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: METHOD FOR PRODUCING ALTERED STARCH FROM POTATO PLANTS



(57) Abstract

Disclosed is a method of producing altered starch from transformed potato plants or their progeny, comprising extracting starch from a potato plant, at least the tubers of which comprise at least an effective portion of a starch branching enzyme (SBE) cDNA sequence operably linked in the antisense orientation to a suitable promoter, such that the level of SBE activity is limited to less than 0.8 units per gram tuber. Also disclosed are potato plants comprising altered starch in accordance with the invention.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

Title: Method for producing altered starch from potato plants

Field of the Invention

This invention relates to a method of obtaining novel types of starch from potato plants, to novel potato plants from which the starch may be obtained, and to vectors for obtaining said plants.

Background of the Invention

Starch is the major form of carbon reserve in plants, constituting 50% or more of the dry weight of many storage organs - e.g. tubers, seeds of cereals. Starch is used in numerous food and industrial applications. In many cases, however, it is necessary to modify the native starches, via chemical or physical means, in order to produce distinct properties to suit particular applications. It would be highly desirable to be able to produce starches with the required properties directly in the plant, thereby removing the need for additional modification. To achieve this via genetic engineering requires knowledge of the metabolic pathway of starch biosynthesis. This includes characterisation of genes and encoded gene products which catalyse the synthesis of starch. Knowledge about the regulation of starch biosynthesis raises the possibility of re-programming biosynthetic pathways to create starches with novel properties that could have new commercial applications.

The commercially useful properties of starch derive from the ability of the native granular form to swell and absorb water upon suitable treatment. Usually heat is required to cause granules to swell in a process known as gelatinisation, which has been defined (W.A. Atwell et al., Cereal Foods World 33, 306-311, 1988) as "....the collapse (disruption) of molecular orders within the starch granule manifested in irreversible changes in properties such as granular swelling, native crystallite melting, loss of birefringence, and starch solubilisation. The point of initial gelatinisation and the range over which it occurs is governed by starch concentration, method of observation, granule type, and heterogeneities

within the granule population under observation". A number of techniques are available for the determination of gelatinisation as induced by heating, a convenient and accurate method being differential scanning calorimetry, which detects the temperature range and enthalpy associated with the collapse of molecular orders within the granule. To obtain accurate and meaningful results, the peak temperature of the endotherm observed by differential scanning calorimetry is usually determined.

The consequence of the collapse of molecular orders within starch granules is that the granules are capable of taking up water in a process known as pasting, which has been defined (W.A. Atwell et al., *Cereal Foods World* 33, 306-311, 1988) as "...the phenomenon following gelatinisation in the dissolution of starch. It involves granular swelling, exudation of molecular components from the granule, and eventually, total disruption of the granules". The best method of evaluating pasting properties is considered to be the viscoamylograph (Atwell et al., 1988) in which the viscosity of a stirred starch suspension is monitored under a defined time/temperature regime. A typical viscoamylograph profile for potato starch is shown in Figure 5, in which the initial rise in viscosity is considered to be due to granule swelling. At a certain point, defined by the viscosity peak, granule swelling is so extensive that the resulting highly expanded structures are susceptible to mechanically-induced fragmentation under the stirring conditions used. With increased heating and holding at 95°C, further reduction in viscosity is observed due to increased fragmentation of swollen granules. This general profile (Figure 5) has previously always been found for native potato starch. In addition to the overall shape of the viscosity response in a viscoamylograph, a convenient quantitative measure is the temperature of initial viscosity development (onset). Figure 2 shows a typical viscosity profile for starch (Kennedy & Cabalda, *Chem. in Britain*, November 1991, 1017-1019), during and after cooking, with a representation of the physical state of the starch granules at various points. The letters A, B, C and D correspond to the stages of viscosity onset (A), maximum viscosity (B), complete dispersion (C) and re-association of molecules (or retrogradation, D).

The properties of potato starch are useful in a variety of both food and non-food (paper, textiles, adhesives etc.) applications. However, for many applications, properties are not

optimum and various chemical and physical modifications well known in the art are undertaken in order to improve useful properties. Two types of property manipulation which would be of use are firstly the controlled alteration of gelatinisation and pasting temperatures and, secondly, starches which do not suffer as much granular fragmentation during pasting as illustrated in Figure 1. Currently the only ways of manipulating the gelatinisation and pasting temperatures of potato starch are by the inclusion of additives such as sugars, polyhydroxy compounds or salts (Evans and Haisman, *Starke* 34, 224-231, 1982) or by extensive physical or chemical pre-treatments (e.g. Stute, *Starke* 44, 205-214, 1992). The reduction of granule fragmentation during pasting can be achieved either by extensive physical pre-treatments (Stute, *Starke* 44, 205-214, 1992) or by chemical cross-linking. Such processes are inconvenient and inefficient. It is therefore desirable to obtain plants which produce starch which intrinsically possesses such advantageous properties.

Starch Biosynthesis

Starch consists of 2 major components: amylose, a linear polymer of alpha, 1-4 linked glucose units; and amylopectin, a branched polymer consisting of an alpha, 1-4 linked glucan backbone with alpha, 1-6 linked branches. The key enzymes in starch biosynthesis are the starch synthases and starch branching enzyme [alpha-1,4-glucan: alpha-1,4-glucan 6-glucosyltransferase, EC 2.4.1.18]. Amylose is synthesized from adenosine 5'-(alpha-D-glucopyranosyl pyrophosphate), or "ADP-glucose", by a starch synthase which is associated with the starch granule: the so-called "granule bound starch synthase" (GBSS). Amylopectin is synthesized from ADP-glucose by the concerted action of a soluble starch synthase (SSS) and starch branching enzyme (SBE). SBE hydrolyses the linear alpha-1-4 glucan chain and rejoins the cleaved portion via an alpha-1-6 linkage to produce a branched structure. The activity of SBE is thus of crucial importance in determining the type, and hence properties, of starch synthesized within plant systems.

Starch Branching Enzyme

In most plant species, SBE occurs in multiple forms (e.g. maize kernels, Boyer & Preiss,

Biochem. Biophys. Res. Commun. 80, 169-175 (1978); sorghum seed, Boyer, Phytochem. 24, 15-18 (1985); rice endosperm, Smyth, Plant Sci. 57, 1-8 (1988); pea embryo, Smith, Planta 175, 270-279 (1988)). However, in potato tuber, only a single form of SBE has so far been identified (Blennow & Johansson, Phytochem. 30, 437-444 (1991)).

Endosperm of maize contains three forms of SBE, namely SBE I, SBE IIa and SBE IIb. The "amylose extender" (ae) mutation causes a large reduction of SBE activity and in particular loss of SBE IIb. This reduction in SBE activity results in a higher ratio of amylose to amylopectin in endosperm starch compared to normal maize (Boyer & Preiss, Biochem. Biophys. Res. Commun. 80, 169-175 (1978)).

In pea embryos, 2 forms of SBE exist. The r (wrinkled) mutant of pea lacks SBE I activity and starch from this source has a higher ratio of amylose to amylopectin than normal peas [Smith, Planta 175, 270-279 (1988)].

In potato, amylose-free mutants have been obtained by X-ray irradiation (Hoverkamp-Hermelink et al., Theor. Appl. Genet. 75, 217-221, 1987) and by transformation with antisense-GBSS constructs (Visser et al., Mol. Gen. Genet. 225, 289-296, 1991). However, no high amylose mutants of potato exist and efforts to produce such via transformation with antisense SBE constructs have, hitherto, been unsuccessful (e.g. DE 41 04782A1). In respect of the latter, Wilmitzer et al., [Proceedings International Symposium on Plant Polymeric Carbohydrates, ed. Meuser, Manners & Siebel (1992) pp 33-39] have, using antisense SBE technology, produced tubers containing only 10-20% SBE activity of control tubers, but: "neither the amylose content of the starch in the tubers of these plants, nor the total starch content of the tubers, was altered" (p.39). Similarly, WO 92/11375 suggests the use of an anti-sense approach to alter the starch content of tubers, but there was no reduction to practice and no data showing success of the approach, which disclosure cannot therefore be considered as enabling.

The present inventors have been able to employ similar techniques to obtain plants with even lower levels of SBE activity than those described by Wilmitzer. Surprisingly, especially in view of Wilmitzer's results, the starch obtained from such plants has

unexpected novel, commercially useful properties.

Summary of the Invention

In a first aspect the invention provides a method of producing altered starch from transformed potato plants or their progeny, the method comprising extracting starch from a potato plant, at least the tubers of which comprise at least an effective portion of a starch branching enzyme (SBE) cDNA sequence operably linked in the antisense orientation to a suitable promoter, such that the level of SBE activity is limited to less than 0.8 units per gram tuber.

A unit of SBE activity is defined below.

It is believed that "antisense" methods are mainly operable by the production of antisense mRNA which hybridises to the sense mRNA, preventing its translation into functional SBE polypeptide (eg. Sheehy et al; 1988 PNAS 85, 8805-8809, Van der Rrol et al; Mol. Gen. Genet. 220, 204-212). Thus, it will be apparent to those skilled in the art that neither a full length SBE cDNA sequence nor a "native" SBE cDNA sequence is essential. Preferably the effective portion of an SBE cDNA sequence comprises at least 2/3 of a full length cDNA sequence, but by simple trial and error, other fragments (smaller or larger) may be found which are functional in limiting the SBE activity to less than 0.8 units per gram tuber. Similarly, the SBE cDNA sequence could be a variant comprising several base mismatches (scattered throughout the sequence or concentrated in a few regions) compared to a native SBE cDNA sequence, yet still give rise to an mRNA molecule capable of inhibiting the translation of mRNA derived from the sense strand of an SBE coding sequence. Such fragments and variants are within the scope of the invention.

It will also be apparent to those skilled in the art that the sequence need not be a cDNA sequence according to the strict sense of the term, in that the sequence used could be an equivalent derived from a genomic SBE encoding sequence, although such genomic sequences will preferably be adapted (e.g. by the removal of intron sequences).

Altered starch produced according to the method of the invention is found to have the following physical properties:

- a) elevated peak temperature of gelatinisation as determined by differential scanning calorimetry (DSC) relative to unaltered starch produced from equivalent non-transformed plants; and
- b) elevated viscosity onset temperature, relative to unaltered starch produced from equivalent non-transformed plants.

The altered starch possesses these qualities *ab initio* as first extracted from the potato plant: the properties are not, for example, acquired by heating in the extraction process.

In a further aspect, the invention thus provides altered starch extracted from transformed potato plants or their progeny having less than 0.8 units SBE activity per gram tuber, the altered starch as extracted preferably having *ab initio* the properties defined above.

The parameters given above are frequently used by those skilled in the art to determine the properties of starch. The Examples below describe particular assay methods by which these parameters may be determined.

The peak temperature of gelatinisation is the temperature at which there is a maximum in the loss of order in granules within a sample of starch in the presence of excess water, as judged by the heat flow required to maintain a constant rate of temperature increase, compared with a sample of water. Preferably the peak temperature of gelatinisation is elevated by at least 2°C, more preferably by at least 5°C, compared to unaltered starch.

For the purposes of the present specification, the viscosity onset temperature is defined as the temperature at which the viscosity of a 10% w/w aqueous starch solution becomes at least 50% greater than the maximum viscosity of the solution at lower temperatures (above 50°C). Viscosity may be measured in arbitrary units (e.g. instrument stirring number units or "SNU"). Preferably the viscosity onset temperature is elevated by at least

3°C, and more preferably by at least 5°C, compared to unaltered starch.

Preferably the altered starch produced from the transformed plants (or the progeny thereof) has a peak temperature of gelatinisation (as determined by differential scanning calorimetry) of at least 71°C and/or a viscosity onset temperature of at least 71°C.

Preferably the plants used in the method comprise a full length SBE cDNA sequence operably linked in the antisense orientation to a suitable promoter.

The altered starch is extracted from potato plants in which the starch branching enzyme (SBE) activity is less than 0.8 units per gram tuber. (A unit of activity is defined for present purposes as the amount of enzyme activity which incorporates into starch 1 micromole of glucose per minute at a temperature of 30°C.)

Preferably the altered starch is extracted from the plant by wet milling of potato tubers.

Preferably the altered starch is obtained from transformed potato plants or their progeny, the tubers of which exhibit less than 10%, and preferably less than 5%, of SBE activity compared to equivalent non-transformed control plants.

In a further aspect, the invention provides a vector for modifying a potato plant so as to cause the plant to be capable of giving rise to tubers having less than 0.8 units SBE activity per gram tuber, the vector comprising at least an effective portion of an SBE cDNA sequence operably linked in the antisense orientation to a suitable promoter.

Preferably the vector comprises a full length SBE CDNA sequence, preferably that of potato SBE, operably linked in the antisense orientation to a suitable promoter. Suitable promoters include the CaMV 35S and the GBSS promoters. In a preferred embodiment the vector comprises a plurality of copies of the CaMV 35S promoter, preferably operably linked in a tandem arrangement.

In another aspect the invention provides a potato plant capable of giving rise to tubers

having less than 0.8 units SBE activity per gram tuber and comprising at least an effective portion of an SBE cDNA sequence operably linked in the antisense orientation to a suitable promoter. Typically, such a plant will have been transformed with an antisense SBE construct, or will be the progeny of such a plant.

Preferably the plant tubers exhibit less than 10%, more preferably less than 5%, of the SBE activity of equivalent non-transformed control plants.

The various aspects of the invention will now be further illustrated by way of example and with reference to the drawings, of which:

Figure 1 shows how the degree of gelatinisation of an unaltered starch sample varies with temperature, as measured by differential scanning calorimetry;

Figure 2 shows the typical viscosity profile of conventional starch during and after cooking, together with representations of the physical state of starch granules at various stages;

Figure 3 shows how the degree of gelatinisation of a sample of altered starch in accordance with the invention varies with temperature as measured by differential scanning calorimetry (DSC);

Figure 4 is a graph of peak temperature of gelatinisation ($^{\circ}\text{C}$) (as measured by DSC) against SBE activity (Units), showing how the two parameters are correlated;

Figure 5 is a graph of viscosity (SNU) against temperature ($^{\circ}\text{C}$) for unaltered starch;

Figure 6 is a graph of viscosity onset temperature ($^{\circ}\text{C}$) against SBE activity (Units), showing how the two parameters are related;

Figure 7 is a graph of viscosity (SNU) against temperature ($^{\circ}\text{C}$) for altered starch in accordance with the invention; and

Figure 8 shows the sequence of a full length potato SBE cDNA clone.

Examples

Example 1 - Construction of Plant Transformation Vectors containing Antisense Starch Branching Enzyme Genes

(a) Construction of Enhanced 35S Antisense Potato Starch Branching Enzyme Plant Transformation Vector

Initially a 1.4 kb EcoRI partial length cDNA for potato starch branching enzyme was purchased from the Agricultural Genetics Company (Cambridge, UK). This cDNA was isolated from a lambda phage library (methylase protected fragments) made from RNA extracted from potato tubers (cv Desiree) using standard techniques (Sambrook, Fritsch & Maniatis, (1989) Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Lab, NY, 2nd Ed). Subsequently a partial cDNA clone of about 2.3kb was isolated containing an additional 0.9 kb of sequence 3' to the original cDNA and including the polyadenylated tail.

Even later a full length clone was obtained and sequenced (shown in Figure 8, Seq ID No. 1), although only partial cDNA clones had been isolated by the time antisense experiments (described below) were conducted. The full length sequence shown in Figure 8 is in reasonably close agreement with the full length sequence of potato SBE disclosed by Poulsen & Kreiberg (1993, Plant Physiol. 102, 1053-1054), although some differences are readily apparent. Other SBE sequences have also been described (Jens Rossmann PhD Thesis, Lebensmitteltechnologie und Biotechnologie der Technischen Universitat Berlin 1992), and again, there are sequence differences with the sequence shown in Figure 8. Nevertheless, in principle, it should prove possible to use sequences derived from, or based on, those disclosed in the prior art to obtain the present invention.

The 5' end of the two partial cDNAs obtained by the inventors had the same EcoRI site (at nucleotides 615-620 in Figure 8). The 3' end of the 2.3kb clone is at nucleotide 3080

of Figure 8 (which includes an EcoRI linker). The 2.3 kb EcoRI fragment was subcloned in an antisense orientation between the duplicated cauliflower mosaic (CaMV) virus 35S promoter (Cabb-JI strain, equivalent to nucleotide 7040 to 7376 duplicated upstream of 7040 to 7433) and the CaMV polyadenylation signal nt 7435-126 (Franck A, Guilley H, Jonard G, Richards R and Hirth L (1980) *Cell* **21**, 285-294) in the vector pJIT 60 (Guerineau et al., (1992) *Plant Mol. Biol.* **18**, 815-818). The promoter-antisense potato starch branching enzyme-polyA fragment was then cloned into the plant transformation vector BIN19 (Bevan M (1984) *Nucl. Acids Res.* **12**, 8711-8721).

(b) Construction of Patatin Promoter Antisense Potato Starch Branching Enzyme Plant Transformation Vector

The 2.3 kb EcoRI fragment (corresponding to about 2/3 of the full length cDNA) of the potato starch branching enzyme was subcloned into the EcoRI site of the pBSSK II plus vector (Stratagene) to create pSJ5. A XhoI (Klenow repaired) SacI fragment containing the SBE fragment from pSJ5 was subcloned into pBI140.5 cut with SmaI and SacI; this places the SBE in an antisense orientation with respect to the promoter. The resulting plasmid was termed pSJ7. For information, pBI140.5 is a BIN19 derivative containing a 3.5 kb Patatin type I promoter (HindIII to DraI of PAT21, Bevan M, Barker R, Goldsborough A, Jarvis M, Ravanagh T & Iturriaga G (1986) *Nucl. Acids Res.* **14**, 4625-4638) and the polyadenylation signal of the nopaline synthase (Bevan M, Barnes W & Chilton M-D. *Nucl. Acids Res.* **11**, 369-385). *E. coli* strain DH5 α was transformed with pSJ7 using standard techniques and the transformant deposited at the National Collections of Industrial and Marine Bacteria, 23 St Machar Drive, Aberdeen AB2 1RY, United Kingdom (date of deposit: 02/12/94; accession number NCIMB 40701).

(c) Transformation of *Agrobacterium tumefaciens*

The plant transformation vectors containing antisense branching enzyme genes were transferred into *A. tumefaciens* (C58/pGV3850) using a direct DNA uptake protocol [An et al., Binary Vectors, In: Plant Molecular Biology Manual (ed. Galvin and Schilperoort) A3 (1988) 1-19] .

Example 2 - Transformation of Potato with Antisense Starch Branching Enzyme Constructs

(a) Stock Cultures

Stock nodal cutting cultures of potato (cv. Desiree) were maintained on Murashige and Skoog basal media (MS) containing 1% sucrose at 22°C in an illuminated culture room (40 μ joules/m²/hr) with a 16h day. Cuttings were taken every three weeks, with 5 plantlets grown in each Magenta vessel to produce nodes with large leaves [Westcott R, Proc. 5th Intl. Cong. Plant Tissue and Cell Culture (1982), ed. Fujiwara, Tokyo]. Establishment of plants into compost was as described by Westcott (1982).

(b) Tuberisation

Tuberisation was achieved by transfer of single nodes to MS media containing 8% sucrose and 2.5 mg/l benzylaminopurine (BAP) and incubating in darkness at 22°C. After tuberisation had proceeded to pea-sized tubers the explants could be transferred to Magenta vessels containing the same media for storage of up to 6 months.

(c) Agrobacterium Infection

Halved *in vitro* tubers were incubated with log phase *A. tumefaciens* cells for 10 min, after which the explant tissue was removed, blotted on filter paper and transferred onto nurse plates. Nurse plates were prepared by plating 2ml *Nicotiana plumbaginifolia* suspension cells (Barfield et al., Plant Cell Reports 4, 104-107 (1985)) onto regeneration media (0.8% Bactoagar, MS salts, 1% sucrose, 0.2 mg/l indole acetic acid (IAA), 5 mg/l zeatin). Explants were incubated under illumination for 2 days before transfer to fresh regeneration media containing 500 mg/l cefotaxime. 5 days later explants were transferred to the same media containing 100 mg/l kanamycin. After 4 weeks (2 transfers) explants were transferred onto expansion media (MS salts, 1% sucrose, 1.0 mg/l gibberellic acid (GA3) containing cefotaxime and kanamycin. After a total of 8 weeks, regenerating shoots were removed and transferred to basal media (MS salts, 1% sucrose) containing cefotaxime and

kanamycin.

(d) Growth of Plants

Rooted regenerants, 1-2cm high, were transferred to compost (50% Levingtons/50% grit) and grown under high illumination ($400 \mu\text{joules}/\text{m}^2/\text{hr}$) at 20°C day and 18°C night with a 16 hr day period. After 10-12 days, plantlets transferred to 3" pots containing Arthur Bowes Universal Compost. After establishment (40 days), four plants from each clone were repotted together in 10" pots with same compost. Day length was reduced to 11 hr after approximately 100 days growth. Tubers were harvested after foliage senescence (approximately 120 days).

Example 3 - Analysis of Transgenic Plants

(a) Southern Analysis

DNA was isolated from leaves of regenerated plants (Dellaporta, Plant Mol. Biol. Reporter 1, 19-21 (1983)), digested with EcoRI, electrophoresed in a 1% agarose gel in TBE buffer, transferred to Genescreen in $20 \times \text{SSC}$ and u.v. cross-linked (Stratalinker, Stratgene). Blots were hybridised to random-prime labelled (Amersham) 2.3 kb EcoRI potato starch branching enzyme fragment in $5 \times \text{SSPE}$ (0.9M NaCl , $50\text{mM NaH}_2\text{PO}_4$, 5mM EDTA), $5 \times \text{Denhardtts}$ solution, 1% SDS, $100 \mu\text{g}/\text{ml}$ denatured salmon sperm DNA at 65°C overnight. Final washing stringency was $0.2 \times \text{SSC}$, 1% SDS at 65°C for 15 min. Positive transformants were identified by hybridising 1.4 and 0.9 kb fragments (endogenous SBE genes produced higher molecular weight hybridising fragments, presumably due to the presence of introns).

(b) Starch Branching Enzyme (SBE) Assay of Transgenic

Tubers

Sample tubers from each plant were taken after harvest, washed and stored at -20°C until assay.

Frozen tubers were crushed in a mortar and pestle in 2 vol. extraction buffer cooled to 4°C. The buffer contained 100 mM 2-amino-2-(hydroxymethyl)-1,3 propanediol (Tris) pH 7.5, 10mM ethylene-diaminetetra-acetic acid (EDTA), 2.5 mM dithiotreitol (DTT), 0.1% (w/v) sodium metabisulphite and 10% (w/v) polyvinyl-polypyrrolidone (PVPP). When completely homogenised the crude homogenate was clarified by centrifuging at 10,000g for 10 minutes. The supernatant was retained for the assay of starch branching enzyme activity.

The standard SBE assay reaction mixture, in a volume of 0.2 ml, was 200 mM 2-(N-morpholino) ethanesulphonic acid (MES) buffer, pH 6.5, 50mM [^{14}C]glucose 1-phosphate (100 nCi), 0.05 mg rabbit phosphorylase A and potato tuber extract. Incubations were performed at 30°C for 60 minutes. Negative controls contained either: (a) no phosphorylase, or (b) the potato tuber extract boiled for 30 minutes to destroy enzyme activity. The reaction was terminated and glucan polymer precipitated by the addition of 1 ml of 75% (v/v) methanol, 1% (w/v) potassium hydroxide (KOH) and then 0.1 ml of glycogen (10 mg ml⁻¹). Insoluble glucan polymer was pelleted by centrifugation and washed with a further 1 ml of methanol/KOH before being redissolved in water and the incorporated radioactivity measured in a Beckman LS 3800 liquid scintillation counter.

Activity was expressed as units. with one unit defined as 1 micromole of glucose incorporated per minute. All measurements were taken during the phase of the assay when the rate of glucose incorporation was linear.

The results are shown in Table 1. For the transgenic plants it can be seen that, relative to control values, SBE activity has been reduced by varying degrees. Several plants have SBE activities less than 0.8U/g tuber (below 10% of average control values).

Starch Branching Enzyme Assays of Transgenic Potato Tuber Extracts

All starch branching enzyme activities were measured in duplicate and mean values taken. At low levels of activity absolute quantitation, via the standard phosphorylase assay, is more difficult because inaccuracies introduced by background activity are proportionally much greater.

POTATO TUBER STARCH BRANCHING ENZYME ACTIVITY		
	PLANT	ACTIVITY (units g ⁻¹ tuber)
<u>CONTROL</u>	58	21.3
	40	18.2
	31	16.6
	29	13.1
	49	13.0
	8	12.7
<u>Pat AS 2/3 Pot</u>	47	2.8
	54	2.4
	69	0.2

POTATO TUBER STARCH BRANCHING ENZYME ACTIVITY		
	PLANT	ACTIVITY (units g ⁻¹ tuber)
<u>2 x 35S AS 2/3 Pot</u>	25	16.9
	5	13.0
	9	13.0
	32	12.5
	16	12.2
	6	11.1
	22	7.7
	23	7.6
	20	6.6
	26	5.5
	34	5.2
	14	4.6
	24	4.6
	61	4.5
	4	4.3
	21	3.9
	19	2.4
	17	2.3
	28	2.3
	18	1.9
	3	1.8
	13	1.4
	10	1.1
	2	1.0
	1	0.7
	53	0.6
	27	0.6
	12	0.5
	15	0.5
	33	0.5
	52	0.5
	11	0.4
	60	0.4
	7	0.4
	72	0.3
	68	0.3
	35	0.2

Example 4 - Analysis of Transgenic Starch Properties

(a) Starch Extraction

Potato tubers were homogenised in water for 2 min in a Waring blender operating at high speed. The homogenate was washed and filtered (initially 2 mm, then 1 mm filters) using approximately 4L of water per 100g of tubers (6 extractions). Washed starch granules were finally extracted with acetone and air dried.

(b) Differential Scanning Calorimetry

The temperature range for the loss of granule order upon heating starches in excess water was determined by differential scanning calorimetry. Starch powders isolated from a range of transgenic potato plants were analysed using the Perkin Elmer DSC 7 instrument. 1-4mg of starch was accurately weighed into an aluminium sample pan, and water added so that the starch concentration was less than 25% w/v, to give a total sample weight of 10-15mg. An empty reference sample pan was used. A heating rate of 10°C/minute was used to heat the test and reference samples from 25°C to 95°C. Data analysis was performed using the instrument software. Examples of results are shown in Figures 1 and 3. A number of temperature parameters can be obtained from such plots, the most accurate being the peak temperature. A difference in peak temperature of 2-3°C is readily determined as shown by comparison of Figure 1 (peak temperature 69.3°C) and Figure 3 (peak temperature 72.0°C).

Starches isolated from potato plants exhibiting a range of starch branching enzyme activities (determined as described in Example 3b) were characterised by differential scanning calorimetry. Peak temperatures are compared with starch branching enzyme activity in Figure 4, from which it appears that levels of enzyme activity less than 0.8U/g of tuber are required for consistent increases in peak temperature.

(c) Viscosity Development

Starches isolated from a range of transgenic potato plants were analysed for viscosity development ('pasting') following the loss of granule order. The instrument used was the Rapid Visco Analyser 3C (Newport Scientific, Sydney, Australia). Starch (2.50g) was weighed into an instrument sample holder, and water (22.50g) added so that the final concentration was 10% w/w starch. Suspensions were equilibrated for 2 minutes at 50°C and heated under standard stirring conditions at 1.5°C minute from 50°C to 95°C, then held at 95°C for 15 minutes. The viscosity developed was measured in instrument stirring number units (SNU). A typical trace obtained is shown in Figure 5. The broad maximum observed as a function of temperature makes the accurate determination of a peak temperature difficult, but the fact that viscosity starts from a very low level and rapidly rises allows an accurate determination of a viscosity onset temperature, defined as the temperature at which viscosity is at least 50% higher than at all lower temperatures above 50°C.

The viscosity onset temperatures for starches isolated from potato plants exhibiting a range of starch branching enzyme activities were determined, with the results shown in Figure 6. These data show that a consistent increase in viscosity onset temperature is found for starches from plants containing less than 0.8U/g of tuber of starch branching enzyme. For those starches which show a higher viscosity onset temperature, other parameters of pasting (e.g. peak temperature) are also higher. This is illustrated by comparison of Figures 5 (onset temperature: 70°C, peak temperature: 82°C) and 7 (onset temperature 75°C, peak temperature: 87°C).

Example 5

Construction of GBSS antisense full length potato starch branching enzyme vector.

The inventors have recently made a further construct comprising a full length potato SBE cDNA in the anti-sense orientation under the control of the GBSS promoter. Details of the construction are given below. No experimental data regarding this construct are yet available.

A full length cDNA clone for potato starch branching enzyme corresponding to nucleotides 91-3114 plus an additional 10 bases at the 3' end (Poulsen, P. & Kreiberg, J.D. *Plant Physiol.* (1993) 102: 1053-1054) was isolated from a potato tuber cDNA library (see above). The cDNA was excised from the plasmid vector by cutting with *SacI* and *XhoI* and inserted in an antisense orientation between the granule bound starch synthase promoter (GBSS) and the nos polyadenylation signal in the BIN 19 based plant transformation vector pPGB121 which had been cut with *SacI* and *SalI*. The GBSS promoter is a 0.8 kb *HindIII* - *NsiI* fragment of the granule bound starch synthase genomic clone LGBSSwt-6; this promoter fragment directs GUS expression in an organ specific manner (up to 3350 fold higher in tubers than in leaves and up to 25 fold higher than the CaMV promoter) (Visser, R G F, Stollte, A. and Jacobsen, E. *Plant Mol. Biol.* (1991) 17:691-699).

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

- (A) NAME: National Starch and Chemical Investment Holding Corporation
- (B) STREET: Suite 27, 501 Silverside Road
- (C) CITY: Wilmington
- (D) STATE: Delaware
- (E) COUNTRY: United States of America
- (F) POSTAL CODE (ZIP): 19809

(ii) TITLE OF INVENTION: Improvements in or Relating to Starch I

(iii) NUMBER OF SEQUENCES: 1

(iv) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 3128 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

GAATTCGGCA CGAGAGCTGA AGCAAAGTAC CATAATTTAA TCAATGGAAA TTAATTTCAA	60
TGTTTTGTCA AAACCCATTC GAGGATCTTT TCCATCTTCC TCACCTAAAG TTTCTTCAGG	120
GGCTTCTAGA AATAAGATAT GTTTTCCTTC TCAACATAGT ACTGGACTGA AGTTTGGATC	180
TCAGGAACGG TCTTGGGATA TTTCTTCCAC CCCAAAATCA AGAGTTAGAA AAGATGAAAG	240
GATGAAGCAC AGTTCAGCTA TTTCCGCTGT TTTGACCGAT GACAATTCGA CAATGGCACC	300
CCTAGAGGAA GATGTCAAGA CTGAAAATAT TGACCTCCTA AATTTGGATC CAACTTTGGA	360

ACCTTATCTA GATCACTTCA GACACAGAAT GAAGAGATAT GTGGATCAGA AAATGCTCAT	420
TGAAAAATAT GAGGGACCCC TTGAGGAATT TGCTCAAGGT TATTTAAAAT TTGGATTCAA	480
CAGGGAAGAT GGTTCATAG TCTATCGTGA ATGGGCTCCT GCTGCTCAGG AAGCAGAAGT	540
TATTGGCGAT TTCAATGGAT GGAACGGTTC TAACCACATG ATGGAGAAGG ACCAGTTTGG	600
TGTTTGAGT ATTAGAATTC CTGATGTTGA CAGTAAGCCA GTCATTCCAC ACAACTCCAG	660
AGTTAAGTTT CGTTTCAAAC ATGGTAATGG AGTGTGGGTA GATCGTATCC CTGCTTGGAT	720
AAAGTATGCC ACTGCAGACG CCACAAAGTT TGCAGCACCA TATGATGGTG TCTACTGGGA	780
CCCACCACCT TCAGAAAGGT ACCACTTCAA ATACCCTCGC CCTCCCAAAC CCCGAGCCCC	840
ACGAATCTAC GAAGCACATG TCGGCATGAG CAGCTCTGAG CCACGTGTAA ATTCGTATCG	900
TGAGTTTGCA GATGATGTTT TACCTCGGAT TAAGGCAAAT AACTATAATA CTGTCCAGTT	960
GATGGCCATA ATGGAACATT CTTACTATGG ATCATTTGGA TATCATGTTA CAAACTTTTT	1020
TGCTGTGAGC AATAGATATG GAAACCCGGA GGACCTAAAG TATCTGATAG ATAAAGCACA	1080
TAGCTTGGGT TTACAGGTTT TGGTGGATGT AGTTCACAGT CATGCAAGCA ATAATGTCAC	1140
TGATGGCCTC AATGGCTTTG ATATTGGCCA AGGTTCTCAA GAATCCTACT TTCATGCTGG	1200
AGAGCGAGGG TACCATAAGT TGTGGGATAG CAGGCTGTTC AACTATGCCA ATTGGGAGGT	1260
TCTTCGTTTC CTTCTTTCCA ACTTGAGGTG GTGGCTAGAA GAGTATAACT TTGACGGATT	1320
TCGATTTGAT GGAATAACTT CTATGCTGTA TGTTTCATCAT GGAATCAATA TGGGATTTAC	1380
AGGAACTAT AATGAGTATT TCAGCGAGGC TACAGATGTT GATGCTGTGG TCTATTTAAT	1440
GTTGGCCAAT AATCTGATTC ACAAGATTTT CCCAGACGCA ACTGTTATTG CCGAAGATGT	1500
TTCTGGTATG CCGGGCCTTA GCCGGCCTGT TTCTGAGGGA GGAATTGGTT TTGATTACCG	1560
CCTGGCAATG GCAATCCCAG ATAAGTGGAT AGATTATTTA AAGAATAAGA ATGATGAAGA	1620
TTGGTCCATG AAGGAAGTAA CATCGAGTTT GACAAATAGG AGATATACAG AGAAGTGTAT	1680
AGCATATGCG GAGAGCCATG ATCAGTCTAT TGTCGGTGAC AAGACCATTG CATTTCTCCT	1740
AATGGACAAA GAGATGTATT CTGGCATGTC TTGCTTGACA GATGCTTCTC CTGTTGTTGA	1800
TCGAGGAATT GCGCTTCACA AGATGATCCA TTTTTCACA ATGGCCTTGG GAGGAGAGGG	1860
GTACCTCAAT TTCATGGGTA ACGAGTTTGG CCATCCTGAG TGGATTGACT TCCCTAGAGA	1920
GGGCAATAAT TGGAGTTATG ACAAATGTAG ACGCCAGTGG AACCTCGCAG ATAGCGAACA	1980
CTTGAGATAC AAGTTTATGA ATGCATTTGA TAGAGCTATG AATTCGCTCG ATGAAAAGTT	2040

20

CTCATTCTC GCATCAGGAA AACAGATAGT AAGCAGCATG GATGATGATA ATAAGGTTGT	2100
TGTGTTTGAA CGTGGTGACC TGGTATTTGT ATTCAACTTC CACCCAAATA ACACATACGA	2160
AGGGTATAAA GTTGGATGTG ACTTGCCAGG GAAGTACAGA GTTGCACTGG GCAGTGATGC	2220
TTGGGAATTT GGTGGCCATG GAAGAGCTGG TCATGATGTT GACCATTTCAT CATCACCAGA	2280
AGGAATACCT GGAGTTCCAG AAACAAATTT CAATGGTCGT CCAAATTCCT TCAAAGTGCT	2340
GTCTCCTGCG CGAACATGTG TGGCTTATTA CAGAGTTGAT GAACGCATGT CAGAAACTGA	2400
AGATTACCAG ACAGACATTT GTAGTGAGCT ACTACCAACA GCCAATATCG AGGAAAGTGA	2460
CGAGAACTT AAAGATTCAT CATCTACAAA TATCAGTACA TCATCTACAA AAAATGCTTA	2520
TTACAGAGTT GATGAACGCA TGTCAGAAGC TGAAGATTAC CAGACAGACA TTTGTAGTGA	2580
GCTACTACCA ACAGCCAATA TCGAGGAGAG TGACGAGAAA CTTGATGATT CATTATCTAC	2640
AAATATCAGT AACATTGGTC AGACTGTTGT AGTTTCTGTT GAGGAGAGAG ACAAGGAACT	2700
TAAAGATTCA CCATCTGTAA GCATCATTAG TGATGCTGTT CCAGCTGAAT GGGCTGATTC	2760
GGATGCAAAC GTCTGGGGTG AGGACTAGTC AGATGATTGA TCGATCCTTC TACGTTGGTG	2820
ATCTTGGTCC GTGCATGATG TCTTCAGGGT GGTAGCATTG ACTGATTGCA TCATAGTTTT	2880
TTTTTTTTTT TAAGTATTTT CTCTATGCAT ATTATTAGCA TCCAATAAAT TTAAGTGGTG	2940
TTGTACATAG AAAAAGTGCA TTTGCATGTA TGTGTTTCTC TGAAATTTTC CCCAGTTTTT	3000
GGTGCTTTGC CTTTGGAGCC AAGTCTCTAT ATGTAATAAG AAAACTAAGA ACAATCACAT	3060
ATATAAAATG TTAGTAGATT ACCATAAAAA AAAAATTAAG AAAAAAAAAA AAAAAGTGA	3120
GGGGGGGC	3128

CLAIMS

1. A method of producing altered starch from transformed potato plants or their progeny, the method comprising extracting starch from a potato plant, at least the tubers of which comprise at least an effective portion of a starch branching enzyme (SBE) cDNA sequence operably linked in the antisense orientation to a suitable promoter, such that the level of SBE activity is limited to less than 0.8 units per gram tuber.
2. A method according to claim 1, wherein the tubers contain at least 2/3 of an SBE cDNA sequence operably linked in the antisense orientation to a suitable promoter.
3. A method according to claim 1 or 2, wherein the starch is extracted from plants, the tubers thereof having less than 10%, and preferably less than 5%, of the SBE activity in equivalent non-transformed plants.
4. A method according to any one of claims 1, 2 or 3, wherein the peak temperature of gelatinisation of the starch so produced is elevated by at least 2°C, preferably by at least 5°C, compared to unaltered starch produced from equivalent non-transformed plants.
5. A method according to any one of the preceding claims, wherein the viscosity onset temperature of the starch so produced is elevated by at least 3°C, and preferably by at least 5°C, compared to unaltered starch produced from equivalent non-transformed plants.
6. A method according to any one of the preceding claims, wherein the peak temperature of gelatinisation (as determined by differential scanning calorimetry) of the starch so produced is at least 71°C.
7. A method according to any one of the preceding claims, wherein the viscosity onset temperature of the starch so produced is at least 71°C.
8. A method according to any one of the preceding claims, comprising wet milling of potato tubers.

9. Altered starch produced by the method of any one of the preceding claims.
10. A vector for modifying a potato plant so as to cause the plant to be capable of giving rise to tubers having less than 0.8 units SBE activity per gram tuber, the vector comprising at least an effective portion of an SBE cDNA sequence operably linked in the antisense orientation to a suitable promoter.
11. A vector according to claim 10, comprising a full length SBE cDNA sequence operably linked in the antisense orientation to a suitable promoter.
12. A vector according to claim 10 or 11, wherein the SBE cDNA sequence is a potato SBE cDNA sequence.
13. A vector according to any one of claims 10, 11 or 12, wherein the vector comprises the CaMV 35S or the granule bound starch synthase (GBSS) promoter.
14. A vector according to any one of claims 10 to 13, comprising a plurality of copies of the CaMV 35S promoter.
15. A vector according to any one of claims 10 to 14, comprising two or more copies of the CaMV 35S promoter operably linked in a tandem arrangement.
16. A vector according to any one of claims 10 to 13, comprising a tuber-specific promoter.
17. A transformed potato plant or the progeny thereof capable of giving rise to tubers having altered starch, and comprising at least an effective portion of an SBE cDNA sequence operably linked in the antisense orientation to a suitable promoter, such that the level of SBE activity is limited to less than 0.8 units per gram tuber.
18. A potato plant according to claim 17, comprising altered starch having a peak temperature of gelatinisation elevated by at least 2°C, preferably by at least 5°C,

compared to unaltered starch produced from equivalent non-transformed plants.

19. A potato plant according to claim 17 or 18, comprising altered starch having a viscosity onset temperature elevated by at least 3°C, and preferably by at least 5°C, compared to unaltered starch produced from equivalent non-transformed plants.

20. A potato plant according to any one of claims 17, 18 or 19, comprising starch having a peak temperature of gelatinisation as determined by differential scanning calorimetry (DSC) of at least 71°C.

21. A potato plant according to any one of claims 17-20, comprising starch having a viscosity onset temperature of at least 71°C.

22. Altered starch extracted from transformed potato plants or their progeny, the plants having less than 0.8 units SBE activity per gram tuber, wherein the starch as extracted has the following physical properties:

a) elevated peak temperature of gelatinisation as determined by differential scanning calorimetry (DSC) relative to unaltered starch extracted from equivalent non-transformed plants; and

b) elevated viscosity onset temperature, relative to unaltered starch extracted from equivalent non-transformed plants.

23. Altered starch according to claim 22, wherein said starch is extracted from plants, the tubers thereof having less than 10%, and preferably less than 5%, SBE activity compared to equivalent non-transformed plants.

24. Altered starch according to claim 22 or 23, wherein the peak temperature of gelatinisation is elevated by at least 2°C, preferably at least 5°C, compared to unaltered starch extracted from equivalent non-transformed plants.

25. Altered starch according to any one of claims 22, 23 or 24, wherein the viscosity onset temperature is elevated by at least 3°C, and preferably at least 5°C, compared to unaltered starch extracted from equivalent non-transformed plants.

1/8





Fig. 1

2/8



CHANGES IN STARCH DURING AND AFTER COOKING

Fig. 2

3/8




Fig. 3




Fig. 4




Fig. 5

6/8




Fig. 6




Fig. 7

10% SUSPENSION

8/8

GAATTCGGCAGGAGCTGAAGCAAAGTACCATAATTTAATCAATGGAAATTAATTTCAA	60
TGTTTTGTCAAAACCCATTGAGGATCTTTTCCATCTTCCCTCACCTAAAGTTTCTTCAGG	120
GGCTTCTAGAAATAAGATATGTTTTCTTCTCAACATAGTACTGGACTGAAGTTTGGATC	180
TCAGGAACGGTCTTGGGATATTTCTTCCACCCCAAAATCAAGAGTTAGAAAAGATGAAAG	240
GATGAAGCACAGTTTCAGCTATTTCCGCTGTTTTGACCGATGACAATTCGACAATGGCACC	300
CCTAGAGGAAGATGTCAAGACTGAAAATATTGACCTCCTAAATTTGGATCCAACCTTTGGA	360
ACCTTATCTAGATCACTTCAGACACAGAATGAAGAGATATGTGGATCAGAAAATGCTCAT	420
TGAAAAATATGAGGGACCCCTTGAGGAATTTGCTCAAGGTTATTTAAAATTTGGATTCAA	480
CAGGGAAGATGGTTGCATAGTCTATCGTGAATGGGCTCCTGCTGCTCAGGAAGCAGAAGT	540
TATTGGCGATTTCAATGGATGGAACGGTCTAACCACATGATGGAGAAGGACCAGTTTGG	600
TGTTTGGAGTATTAGAATTCCTGATGTTGACAGTAAGCCAGTCATTCCACACAACCTCCAG	660
AGTTAAGTTTCGTTTTCAAACATGGTAATGGAGTGTGGGTAGATCGTATCCCTGCTTGGAT	720
AAAGTATGCCACTGCAGACGCCACAAAGTTTGACAGCACCATATGATGGTGTCTACTGGGA	780
CCCACCACCTTCAGAAAGGTACCACTTCAAATACCTCGCCCTCCCAAACCCCGAGCCCC	840
ACGAATCTACGAAGCACATGTGGCATGAGCAGCTCTGAGCCACGTGTAAATTCGTATCG	900
TGAGTTTGCAGATGATGTTTTACCTCGGATTAAGGCAAATAACTATAATACTGTCCAGTT	960
GATGGCCATAATGGAACATTCTTACTATGGATCATTTGGATATCATGTTACAAAATTTTT	1020
TGCTGTGAGCAATAGATATGGAACCCGGAGGACCTAAAGTATCTGATAGATAAAGCACA	1080
TAGCTTGGGTTTACAGGTTCTGGTGGATGTAGTTCACAGTCATGCAAGCAATAATGTCAC	1140
TGATGGCCTCAATGGCTTTGATATTGGCCAAGGTTCTCAAGAATCCTACTTTTCATGCTGG	1200
AGAGCGAGGGTACCATAAGTTGTGGGATAGCAGGCTGTTCAACTATGCCAATTGGGAGGT	1260
TCTTCGTTTCTTCTTCCAACCTTGAGGTGGTGGCTAGAAGAGTATAACTTTGACGGATT	1320
TGATTTGATGGAATAACTTCTATGCTGTATGTTTCATCATGGAATCAATATGGGATTTAC	1380
AGGAAACTATAATGAGTATTTTCAGCGAGGCTACAGATGTTGATGCTGTGGTCTATTTAAT	1440
GTTGGCCAATAATCTGATTCACAAGATTTTCCAGACGCAACTGTTATTGCCGAAGATGT	1500
TTCTGGTATGCCGGGCCCTTAGCCGGCCTGTTTTCTGAGGGAGGAATTGGTTTTGATTACCG	1560
CCTGGCAATGGCAATCCAGATAAGTGGATAGATTATTTAAAGAATAAGAATGATGAAGA	1620
TTGGTCCATGAAGGAAGTAACATCGAGTTTGACAAATAGGAGATATACAGAGAAGTGAT	1680
AGCATATGCCGAGAGCCATGATCAGTCTATTGTGCGGTGACAAGACCATTGCATTTCTCCT	1740
AATGGACAAAGAGATGTATTCTGGCATGTCTTGCTTGACAGATGCTTCTCCTGTTGTTGA	1800
TCGAGGAATTGCGCTTCAAGATGATCCATTTTTTCACAATGGCCTTGGGAGGAGAGGG	1860
GTACCTCAATTTTCATGGGTAAACGAGTTTGGCCATCCTGAGTGGATTGACTTCCCTAGAGA	1920
GGGCAATAAATTGGAGTTATGACAAATGTAGACGCCAGTGGAACCTCGCAGATAGCGAACA	1980
CTTGAGATACAAGTTTATGAATGCATTTGATAGAGCTATGAATTCGCTCGATGAAAAGTT	2040
CTCATTTCTCGCATCAGGAAAACAGATAGTAAGCAGCATGGATGATGATAATAAGGTTGT	2100
TGTGTTTGAACGTGGTGACCTGGTATTTGTATTCAACTTCCACCCAAATAACACATACGA	2160
AGGGTATAAAGTTGGATGTGACTTGCCAGGGAAGTACAGAGTTGCACTGGGCAGTGATGC	2220
TTGGGAATTTGGTGGCCATGGAAGAGCTGGTCATGATGTTGACCATTTACATCACCAGA	2280
AGGAATACCTGGAGTTCCAGAAACAAATTTCAATGGTCTGTCCTTCAAAAGTGCT	2340
GTCTCCTGCGCGAACATGTGTGGCTTATTACAGAGTTGATGAACGCATGTCAGAACTGA	2400
AGATTACCAGACAGACATTTGTAGTGAGCTACTACCAACAGCCAATATCGAGGAAAGTGA	2460
CGAGAACTTAAAGATTCATCATCTACAAATATCAGTACATCATCTACAAAAAATGCTTA	2520
TTACAGAGTTGATGAACGCATGTGAGAAGCTGAAGATTACCAGACAGACATTTGTAGTGA	2580
GCTACTACCAACAGCCAATATCGAGGAGAGTGACGAGAAAACCTTGATGATTCAATTATCTAC	2640
AAATATCAGTAACATTGGTCAGACTGTTGTAGTTTTCTGTTGAGGAGAGAGACAAGGAACT	2700
TAAAGATTCACCATCTGTAAGCATCATTAGTGATGCTGTTCCAGCTGAATGGGCTGATTC	2760
GGATGCAAACGCTCTGGGGTGGGACTAGTCAGATGATTGATCGATCCTTCTACGTTGGTG	2820
ATCTTGGTCCGTGCATGATGCTTTCAGGGTGGTAGCATTGACTGATTGCATCATAGTTTT	2880
TTTTTTTTTTAAGTATTTCTCTATGCATATTATTAGCATCCAATAAAATTTACTGGTTG	2940
TTGTACATAGAAAAAGTGCATTTGCATGTATGTGTTTCTTGAAATTTTCCCAGTTTTT	3000
GGTGCTTTGCCTTTGGAGCCAAGTCTCTATATGTAATAAGAAAACTAAGAACAATCACAT	3060
ATATAAAATGTTAGTAGATTACCATAAAAAAAAAAATTAATAAAAAAAAAAAAAAAAAA	3120
GGGGGGGC	3128

Fig. 8

SUBSTITUTE SHEET (RULE 26)

INTERNATIONAL SEARCH REPORT

International Application No

PLT/GB 95/00634

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/82 C12N15/11 A01H5/00 C08B30/14

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO-A-92 14827 (INSTITUT FÜR GENBIOLOGISCHE FORSCHUNG BERLIN GMBH) 3 September 1992 *whole document* & DE-A-41 04 782 cited in the application ---	1,9-13, 16-18
X	Abstracts VIIth International Congress on Plant Tissue and Cell Culture, Amsterdam, June 24-29, 1990, abstract no. AS-28, F.R. van der Leij et al. "Expression of the gene encoding granule-bound starch synthase after introduction ..." ---	9,22-25
X	WO-A-92 11375 (AMYLOGENE HB) 9 July 1992 cited in the application *whole document* ---	1,9-18
	--- -/--	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *A* document member of the same patent family

Date of the actual completion of the international search

11 July 1995

Date of mailing of the international search report

2 9. 08. 95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Yeats, S

INTERNATIONAL SEARCH REPORT

International Application No
PLT/GB 95/00634

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>PROC. INTERNAT. SYMP. PLANT POLYMERIC CARBOHYDRATES, 1992 pages 33-39, L. WILLMITZER ET AL.; 'Starch synthesis in transgenic plants' cited in the application *page 37, last paragraph - end of page 38*</p>	1
A	<p>PLANT PHYSIOL., vol. 102, 1993 pages 1053-1054, P. POULSEN AND J.D. KREIBERG; 'Starch branching enzyme cDNA from Solanum tuberosum' cited in the application *whole document*</p>	1
A	<p>FEBS LETT., vol. 332, 1993 pages 132-138, J. KHOSHNOODI ET AL.; 'Characterization of the 97 and 103 kDa forms of starch branching enzyme from potato tubers' *whole document*</p>	1
A	<p>MOLEC. GEN. GENET., vol. 225, 1991 pages 289-296, R.G.F. VISSER ET AL.; 'Inhibition of the expression of the gene for granule-bound starch synthase in potato by antisense constructs' cited in the application *abstract; discussion*</p>	1

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 95/00634

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9214827	03-09-92	DE-A- 4104782 AU-A- 1226592 EP-A- 0571427 HU-A- 65740	20-08-92 15-09-92 01-12-93 28-07-94
WO-A-9211375	09-07-92	SE-B- 467160 AU-A- 9109791 EP-A- 0563201 SE-A- 9004095	01-06-92 22-07-92 06-10-93 01-06-92

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.